Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.





J. M. THORBURN & Co.'s

Annual Descriptive Catalogue

VEGETABLE AND AGRICULTURAL SEEDS.

Garden, Field, Fruit, &c., Seeds.

THE LARGEST COLLECTION TO BE FOUND IN THE WORLD.

EMBRACING EVERY

STANDARD AND IMPROVED VARIETY:

TESTED NOVELTIES,

BOTH OF DOMESTIC AND OF FOREIGN ORIGIN, THAT ARE SUITED TO THE CLIMATE OF THE UNITED STATES.

No. 15 JOHN STREET.

Entered according to Act of Congress, in the year 1870, by

JAMES M. THORBURN & CO.,

In the Clerk's Office of the District Court of the United States for the Southern District



NOTICE!!

Our Customers at a distance will much oblique us by sending their orders EARLY. The press of our business is so great, as the sowing season approaches, that some delay in filling orders is unavoidable. Our friends, who send early orders, secure thereby prompt attention. Address, J. M. Thorburn & Co., 15 John STREET, NEW YORK.

Orders should be written apart from the rest of the Letter. Always write the NAME of the variety of Seeds wanted, and not merely the number in the Catalogue.

Be particular to give your NAME and ADDRESS IN FULL with every order, to enable us to forward it promptly and correctly.

Customers are recommended to try the vitality of our Seeds as soon as received. this can readily be done with many Seeds by sowing a few in a flower-pot put in a warm place and watered daily.

We indicate, at the foot of each variety, the most approved for early and late growth, and for any special purpose. These remarks are varied as each year's experience, or the increase of varietics may render necessary, and as we prepare these recommendations with much care they may be fully relied on.

The Directions for Culture are prepared for the neighborhood of New York City. Allowance must be made for more Northerly or Southerly localities as to the time of sowing, etc.

Families residing at a distance may rely upon us to make a proper selection of Seeds for the year, if they think fit, informing us of the size of their garden, or the amount to which it is wished the order should extend.

Our stock of all Seeds for the present year is fully up to our usual supply, both as respects quality and purity, and they may, therefore, be relied upon to give entire satisfaction.

Unknown correspondents sending orders for Seeds, etc., will please send Post Office money order, with order.

SEEDS MAY BE SENT BY MAIL

TO ANY PART OF THE UNITED STATES, AT THE FOLLOWING RATES:

Packag	ges of 4	ounces,	or under	, in	we	ight		 	2	cents.
66	over 4	ounces,	and not	over	8	ounce	S	 	4	66
46	" 8	6.6	66	66	12	66		 	6	44
ei.	" 12	"	"	66	16	44		 	8	66

The maximum weight of Packages which may be sent through the Mails at these Rates is limited to Four Pounds.

RULES

FOR THE

Cultivation of Vegetables.

Both in weight and quality, the crop of all Vegetables to be as large as the quantity and nature of the ground sown permits, will be found to increase in proportion as sufficient space is afforded for the proper tillage of the ground during its growth, and for the admission of the requisite influence and circulation of light and air. In the Directions for Culture appended to the various kinds of Vegetables in the following Catalo ue, the minimum space required is pointed out. As Gardens, in some instances are small, in preparing directions for general use, regard has been had to the due economy of ground, which is the reason that the minimum scale of space has been adopted in the following Catalogue.

The greatest impediment to success in the production of fine Vegetables, next to that pointed out in the preceding paragraph, perhaps, is want of due attention to the proper tillage of the ground. Nothing can supersede the advantages accruing from deep culture. When the plow is used, the sub-soil plow should follow it; and all land, whether heavy or light (except a sub-soil of pure gravel,) will be benefited by it. When the spade is used the ground should be dug at least sixteen inches deep.

Good drainage is of equal importance.

It is best to sow and cultivate ALL crops in drills and rows, in preference to broadcast. The crop is tilled at less expense, both of time and trouble. The hoe, during the growing season, should be continually at work; and, in direct proportion to the complete eradication of the weeds, and to the keeping of the surface of the ground open and loose, will be the weight and quality of the crop.

The rotation of crops should be attended to. Manure should be given to crops above ground, as the Cabbage tribe, Peas, the Potato (for that is not a root.) and similar vegetables. These should be followed by 100t-crops, as Carrots, Parsnips, &c., with less manure.

Those who are experienced cultivators have no need of these suggestions; but few, who are not so, will be convinced of their importance, until by direct experiment they are satisfied of their value and truth.

We take this opportunity to call attention to the advantage as regards the early maturity of many Vegetables to be derived from starting them in a hotbed in Spring. The trouble is not great and the time saved is valuable.

To PREPARE A HOT-BED for this purpose, a quantity of stable-manure should be well shaken up in a heap, about the beginning or middle of February; after remaining three days, it should again be shaken up and turned over, and after an interval of three days repeat the same process. Let the manure remain for three or four days more in the heap, and then form the bed of any dimensions that may be requisit, according to size or number of frames to be used, making the bed about a foot wider than the frame all around. The bed, when made, should not be less than three feet high, the situation for it should be dry underneath, sheltered from the North as much as possible and fully exposed to the Sun. In making up the bed, beat the manure well down with the fork-Place about eight inches of light rich, soil in the frame after it is placed on the bed, so that the surface of the soil may be from six to eight inches from the top of the frame. After the bed has stood with the lights on, but raised a few inches to let the steam escape, for a few days, sow the seed of Cabbage, Tomatoes, Cauliflower, Egg Plant, Celery, Lettuce, Radish, &c., &c. When the Seeds make their appearance above ground, give some air, day and night, in moderate weather, by raising the lights two or three inches, - and as the heat of the bed diminishes, increase the supply of air, to prevent the plants being drawn up. In severe weather, the frame and lights should be covered at night with mats or straw. In high wind, protect the side of the bed also in the same way, or the current of air through will extinguish the heat of the bed. If made early in the season, water only will be required when the earth in the frame becomes quite dry; as the Plants increase in size, the supply of water may be increased. In frosty or cold weather the water should be made milk warm before it is used; this is important.

The under-mentioned Catalogues are Published during the year. all of which will be mailed free on application.

Catalogue of VEGETABLE, AGRICULTURAL, &c., SEEDS, ready in January.

Catalogue of FLOWER SEEDS, containing Directions for their Cultivation, with over 1,000 Varieties, ready in February.

Catalogue of TREE and SHRUB SEEDS, with Directions for their Cultivation, containing over 300 Varieties, ready in February.

Catalogue of FRENCH HYBRID GLADIOLUS, and other Spring Bulbs, ready in February.

Catalogue of DUTCH BULBOUS ROOTS, with Directions for their Management, ready in September.

Catalogue and Almanac, with Directions for the Cultivation of Vegetables in the SOUTHERN STATES, ready in November.

January, May, August and October.

J. M. THORBURN & Co.'s

ANNUAL CATALOGUE

- OF -

Kitchen Garden Seeds,

Etc., etc., etc.

! ! ** See end of this Catalogue for complete assortments of VEGETABLE SEEDS, of approved and choice sorts, for from \$5 to \$25.

Please order by NAME and NOT by the numbers, to facilitate the early execution of Orders.

ARTICHOKE-ARTICHAUT-Alcachofa-Artichode.

French Green Globe, \$\Pi\$ oz., 50 cents; \$\Pi\$ fb., \$5.00. Large Paris, very hardy. \$\Pi\$ oz., \$1.50.

Gulture.—Select deep, rich, light loam, not liable to retain much water in Winter. Give plenty of well-rotted manure. Plant the seed the beginning of May, in an open situation, and keep clear of weeds; if dry weather, water frequently. The seedlings should be left in rows three and a half feet apart, and three feet distant in the rows. When cold weather approaches they should be covered with coarse litter, and every Spring receive a good dressing of well-rotted manure, well worked in with a fork. The offsets should be taken off, and the strongest used for new beds. A bed will last five or six years.

ASPARAGUS-ASPERGE-Esparrago-Spargel.

Giant, Poz, 10 cts. Pho, 60 cts. Conover's Colossal Ppaper of 2 oz., 50cts

ASPARAGUS ROOTS.

Giant, \$1000, \$8; \$100, \$1. Conover's Colossal, \$1000, \$25; \$100, \$3.00

Culture.—The seeds may be sown in the Fall, or as early as the season will permit in the Spring, in drills one foot apart. Press the earth well down with a roller or board. Keep free from weeds, and in the Spring of the second year transplant into permanent beds, which can be made as follows: The ground should not be too wet, nor too strong, and should have a large supply of well-rotted manure, several inches thick, laid on, and then regularly trenched eighteen inches deep, mixing well the manure. The beds should be four and a half feet wide, with walks two feet wide. Four rows of plants should be put in each bed, and the plants twelve inches distant in the rows. The crown of each plant should be about six inches below the level of the bed. Every Fall cut the stalks down; afterwards giving a good dressing of manure, which turn under in the Spring without disturbing the roots. A slight dressing of salt in early Spring is very beneficial. Conover's variety requires more space.

ENGLISH DWARF BEANS-Feve DE MARAIS-Haba-Große Bohnen.

 1. Early Mazagan
 3. Sword Long Pod
 9 qt. 40

 2. Broad Windsor
 40 4. Green Nonpareil
 40 40

Kinds.—No. 1 is the earliest, and Nos. 2 and 3 are the best for main crop.

Culture.—To do well, Broad Beans should be planted as early as possible after the ground is open, in order to get them into pod before the heat of Summer. When in pod, break off the tops of the plant to check growth. Plant and hoe same as for Dwarf Beans. A strong soil suits them best.

DWARF, or SNAP BEARS-HARICOT-Prijoles-Arup-Belinen.

	A bus. A qt.		H	bus.	e qt.
1. Early Snap Shorts.	\$6 00 39	9. Early Rachel	\$	7 00	40
2. Early Turtle Soup.	6 00 30	10. Early Yellow Six			
3. Early Valentine	7 00 40	11 Long do			
4. Early Mohawk		12. Black Wax (fine)			
5. Early China		13. Refugee, or 100			
6. Early Dun Colored	7 00 40	14. Large White Kid			
7. Early Marrowfat (Royal Dwarf.		6 00	30
8. Small White Rice		J			

Kinds.—Nos. 3 and 10 are the earliest; No. 4 the most hardy; Nos. 1, 12 and 13 are very productive, and best for main crop; No. 13 also for late, and the best for pickling; No. 12 is a new Dwarf Wax variety with white transparent pods

Culture.—About the first of May select a warm dry, sleltered spot; dig and manure slightly, make drills an inch deep and two feet apart. Drop the Beans two inches apart in the drill, and cover not more than an inch deep, keep hood when not wet, and the weeds killed; and plant every two weeks for a succession.

POLE, OF RUNNING BEANS—HARICOT-RAMES -Frijole de bejuco— Stangen-Bohnen.

10. French Soisson
11. German Wax (black) " 75
12. White Wax " 75
13. Algiers Wax (black) "1 00
14. Giant Wax (Red) "1 00
15. New Marshall Pole "1 00
16. New Blue Podded (fine). "100
17. Yellow Runners " 1 00
18. French YardLong Runners " 1 00
19. Yard Long # paper 25

Kinds.—Nos. 1, 3, 11 and 18 are best for Snaps.—and Nos. 6, 7 and 10 the best for Shell Beans, and most valuable for main or p.

Culture.—If warm, dry weather, plant about the middle of May for an early crop; and about the first of June plant Limas and the other sorts for the principal crop. Plant in hills, about four feet apart, and not more than two inches deep. Manure freely

REET-BETTERAVE-Remolacha-Runfelrüben.

₽ lt. ₽ oz.	
1. Extra Early Turnip \$1 50 2)	9 Swiss Chard (for greens)\$1 25 15
2. Early Bassano 1 50 20	10. Whyte's Dark Blood 15
	11. Variegated for garnishing). 3 00 30
4. New Dark Egyptian do 50	12. Long Red Mangel Wurtzel 60 10
5. Early Yellow Turnip 1 25 15	13. Yellow Globe Wurtzel 75 10
6. Pine Apple 3.1	14 White Sugar 60 10
7. Carter's St. Osyth 3 00 30	15. Yellow Sugar 75 10
8. Long Smooth Blood 1 0) 15	

Kinds.—Nos. 1 and 2 are the earlie t and best if used when young; No. 3 for general and Winter use; No. 8 for late and Winter crop; Nos from 12 to 15 for cattle feeding.

Culture.—Sow the seed in April for early, and for a succession till last of June; about the middle of May, for general crop. The soil should be a deep, rich, light leam; dry, rather than moist. For the long kinds, trench eighteen inches deep. The rows should be eighteen inches apart, and the plants thinned to eight inches in the rows. Stir the soil often, to kill the weeds, and the plants will make larger roots. As soon as the first hard frost occurs, take up the roots and store away in dry sand.

BORAGE—BOURRACHE—Borraja—Borretch.

Kinds.—The young leaves of this Annual, which smells somewhat like a Cucumber, are used as Salads or boiled as Spinach. Should be sown at intervals, as the leaves are palatable only when young.

Culture.—Sow in Spring, in light soil, and transplant in shallow drills a foot apart when of about six weeks' growth.

BROCOLI-BROCOLI-Broculi-Spargelfehl.

1	. White Sprouting	. # oz.	50 [5. Early Purple Cape (fine) \$\P\$ oz. 6	0
				6. White Cape or Cauliflower " 7	
				7. Sulphur-Colored " 5	
				8. Chappel's Cream " 5	
				adapted to the climate, and the only	

ones that succeed generally. We recommend No. 5 as the most certain.

Culture.—Sow early sorts as soon as the ground can be worked in the Spring in shallow drills, drawn three or four inches apart. Plant out two feet apart each way, when plants are four inches high. Cultivate the same as Cabbage;

rich soil. For a succession, sow one and two months later, taking care to shade the drills.

BRUSSELS SPROUTS-Chou de Bruxelles-Berza de Brusela-Rejentell

30 cents \$2 ounce; \$3 00 \$7 lb.
Improved Dwarf, \$3 ounce 50 cents.

Culture.—Sow in May, in the same manner as Brocoli and transplant in July. They become very tender when touched by frost. Often grow four feet high, and form numerous small heads like Cabbage.

BURNET-PIMPERNELLS DE JARDIN-Pimpernella-Pimpernell.

Garden, 20 cents Pounce,

Culture.—Sow early in Spring, half an inch deep, thinly in rows. Manure freely, every year.

CABBAGE-CHOU-Berza de repollo-Rohl.

CHEST CHOC IS	or sa ac reporte diego.
₩ lt. P oz.	₩ lb.]₩ oz.
1. King of Dwarf\$5 00 50	17. Small Early Ulm Savoy.\$4 00 40
'2. Little Pixie (fine flavor) 4 00 40	18. Early London Market . 3 00 30
3. Earliest Dwarf 3 00 30	19. Early Sugar Loaf 3 00 30
4. Early Vanack 3 00 30	20. Early 0x-Heart (true) . 3 00 30
5. Early Spotsboro' 3 00 30	21. Large Early York 2 00 20
6. Early Victoria 3 00 30	22. Large Late Bergen 3 00 30
7. Early Nonpareil 3 00 30	23. Large Late Drumhead. 3 00 30
8. Early York 2 00 20	24. Large Fine Flat Dutch. 3 03 30
9. Early Salisbury Dwarf. 3 00 30	25. Marblehead Mammoth. 8 00 75
10. Early Dutch 2 00 20	26. Large Late Green Glaz'd 4 00 40
11. Early Wakefield 5 00 50	27. Late Drumhead Savoy. 4 00 40
12. Early Winningstadt 3 00 30	28. Late Globe Savoy 2 00 20
13. Heale's Early Imperial 3 00 - 30	29. Red (for pickling) 4 00 40
14. Early Battersea 3 00 30	30. Thousand Headed 3 00 30
15. Early Drumhead 3 00 30	31. Late Solid H'd Brunsw'k 3 00 30
16. Early Dwarf Savoy 3 00 30	

Kinds.—Nos. 1, 2, 3, 8, 16 and 17, are the earliest; Nos. 10, 12 and 20 are the best for second early. We particularly recommend Nos. 12 and 20. Nos. 22 28 and 24 are the well-known Winter varieties. No. 25 will be found equally worthy of cultivation. No. 24 is the best for main crop. No. 27 is the best for family use.

Culture.—Sow the early sorts in the Autumn and protect by a cold frame during the Winter, transplanting early in Spring; or in hot-beds during February and March, and transplant middle of April; or sow the seed in open ground, as early in the Spring as the ground can be worked. Sow, for later crop, in April and May. They should be sown in shallow drills, four to six inches apart. When the plants are six inches high, transplant into richly manured ground; the early kinds two feet apart, the later kinds, for Winter use, three feet apart. The ground must be deeply loosened and worked thoroughly to grow large and good heads. Hoe often, to kill weeds, and draw some earth ap to the stems. If troubled by the Cabbage flea (Hallica), sprinkle the plants with soot, wood-ashes, lime, or Scotch snuff; either of the above will drive them away.

CARDOON - CARDON - Cardo - Carbon.

Large Solid, 50 cents # ounce.

Culture.—Sow early in Spring, in rows, where they are to stand, and thin them to one foot apart; when full size, bind plants together with straw or matting, and earth up like Celery to bleach.

CARROT-CAROTTE-Zanahoria-Diöhren.

	P lb. Poz.	₹	tb.	Pos.
1.	Extra Early Forcing \$2 00 20	5. Altringham\$	1 50	15
	Early Horn 1 50 15			
	Half Long Red 1 25 15			
	Long Orange 1 25 15			

Rinds.—For the earliest, and for the latest crop. No. 2 is best; Nos. 4, 5 and 7 are the best for main crop; No. 4 is particularly recommended; No. 1 is the best for forcing.

Cuiture.—Sow No. 2 as early as ground can be worked; or, for late crop, until latter part of July; for main crop, sow from the middle of May to first of July. Thin out early crop to five inches in row; main crop to six or seven inches; the rows ten inches apart for early crop; fourteen, for main crop. Hoe often and deeply between rows. Soil, light sandy loam, richly manured and deeply dug.

CAULIFLOWER - CHOU-FLEUR - Coliftor - Blumentohl.

1.	Extra Early Erfurt	P pk	1. 25	6. Early Walcheren Poz.	75
				7. Large White French "	75
3.	Early Paris (fine)		1 50	8. Large Late London "	75
4.	Thorburn's Nonpareil	66	1 25	9. Large Asiatic (true) "	75
5.	Early London	66	75	10. Lenormands (Short Stalk). " 2	00

Kin Is.—Nos. 1, 2 and 3 will be found the flaest varieties for early; No. 1 also for forcing, and No. 4 for late. No. 4 is particularly recommended for general crop; No. 10 is the largest in cultivation.

Culture.—Sow in light, rich soil, about the 20th of August to the middle of September, to transplant into cold frames and protect though the Winter, and transplant early in Spring. to head in June; or sow in hot-bed in February; transplant the latter part of Marchinto another frame, and plant out early in May to head at the same time. Plant three feet by two. For Autumn, or main crop, sow in open situation in May, and transplant in July. Soil must be rich and deep, and, if dry weather, water liberally. As the flower heads appear, the large leaves should be broken down over them, to keep the sun and rain from injuring them.

CELERY-CELERI-Apio-Sellerie.

₩ lb. P (Z.	
1. Incomparable Dwf. White \$5 00 50	4. Dickson's Mammoth White 4 00 40
2. " Crimson 5 00 50	5. Dwarf Self-Blanching 3 00 30
	6. Celeriac or Turnin Root'd 3 00 30

Kinds.—Nos. 1, 2, 3 and 4 are considered the best sorts. We recommend Nos. 1 and 2 for early, and No. 3 for main crop.

Culture. - Sow early in Spring, in a seed-bed half an inch deep, at any time after night frosts are gone, or on a slight hot-bed, if w'shed early; thin out the

plants, when up, to an inch apart, or transplant into a second bed three inches apart. When strong plants, dig trenches four feet apart and fifteen inches wide; dig each trench a moderate spade deep, put three inches of well-rotted dung all over the bottom of each trench, then pare the sides, and dig the dung and parings with an inch or two of loose mould at the bottom, incorporating all well together, and put in the plants in two rows six inches apart, and eight inches apart in rows. Give a good watering after planting, earth up as the plants make growth, and keep the earth between the trenches loose and open with the spade. For the Dwarf variety trench but half the depth. The Turnip-Rooted variety does not require trenches at all; ordinary garden culture is all that is necessary.

CHERVIL—Centeun.—Perifodo — Benfenlaudy. 25 cents % ounce; \$2 50 % lb.

Culture.—Cultivate and use like Parsley. Sow at any time in the Spring, in shallow drills, one foot apart.

TIBLE US CLEAVIL. - Common or our Dumesum. 30 cents #8 onnce.

Culture.—Its treatment is much like the Carrot, except that it should be sown in August and September, in good substantial soil, watering in dry weather, and protecting from slugs by a slight surface dressing of lime or clean ashes. The roots are ripened by July, and can be preserved until April, if occasionally examined to prevent premature growth. They are perfectly hardy, but require a slight covering of litter in the Winter.

COLLARDS-Cnou-Cabu-Blätterfohl.

English, 20 cents P oz.; \$2 00 P lb. Georgia Grown, 40cts. P oz.; \$1 P lb.

Culture.—Sow seeds as for Cabbage, in June, July and August, for succession Transplant when a month old, in rows a foot apart each way, and hoo frequently.

CORN-Mais-Maiz- Mais.

FOR GARDEN CULTURE.

1.	Early Dwarf Sugar	H e	ar. 5	cents;	# qu	art, 40	cents.
	Crosby's Early Sugar		10	6 6	46	50	6.6
	Early Eight-Rowed Sugar	64	8	4.6	44	30	44
	Late Eight-Rowed	66	8	6.	4.6	30	44
	Twelve-Rowed Sugar	6.0	8	66	6.6	30	4.4
	Stowell's Evergreen Sugar	44	8	44	44	30	44
	Trimbles Sugar	44	8	66	66	40	44
	Black Sugar	44	8	. 6	44	40	- 44
	Early Burlington	44	5	44	66	30	4.4
	Early Tuscarora	66	8	44	4.6	30	44
	Japanese Striped Maize	44	10	16	44	1 25	44

Minds.—Nos. 1 and 2 are best for very earliest; No. 1 grows but three feet high; Nos. 3 and 5 for a second or general crop; Nos. 6 and 7 for late; No. 11 is the beautiful striped-leaved v. riety, which is also a fine eating sort.

CHITUTE.—Plant in hills three feet apart cach way, and three or four in a hill; hoe often and draw up soil to the stems; break off side-shoots. Make the ground rich with well-rotted manure.

CORN-FOR FIELD CULTURE.

1.	Early (Canad	la					 	0.0	 	 	 	 	 10	F2.	#	100,	32	50
2.	Early (Conne	ectic	ut	Va	ile:	/	 		 	 	 	 		44		+6	Z	50
3.	Large	Eight	-Ro	we	11	Vhi	te.	 		 	 	 	 		66		4.6	4	00
4.	Eight-	Rowe	dY	ello	W	Fiir	ıt.	 		 	 	 	 		66		6.6	4	00
õ.	Yellow	Dutto	n					 		 	 	 	 		46		66	4	00
6.	Valilie	Dutto	n					 		 							- 6	8	- ()
Pr E -	Orang	e Fii	nt					 		 	 	 	 		6.6		66	-	0)
8.	Pop (o	r Chi	cken) C	01	j.,		 		 	 	 	 		44		bushel,	2	00
																	nded;		
ani	17 are	fine 1	vari	etie	3.														

CORN SALAD, or PETTICUS—Macur—Mache, 6 Valerianilla—

15 cents \$ ounce; \$1 25 \$ ib,

New Striped-Leaved, Pounce 50 cents.

Culture.—Sow latter part of August, or early in September, in drills a quarter of an inch dasp and six inches apart. If dry weather, trend in the seed lightly. Keep down weeds with hoc. Just before the first of Winter, cover thinly with clean straw or leaves.

CRESS-CRESSON-Berro 6 Mastuerzo-Greffe.

1. Curied (or Pepper Grass) \$0 00 10 3. Broad-Leaved Winter....? oz. 30

CUCUMBER-Concomber-Pepino-Gurfenterner.

	P lb. 3	2 oz.	P tb. P cz.
1.	Early Russian\$2 00	20	5. Long Green\$1 50 20
2.	Early Short Green 1 25	15	6. Long White Turkey 4 00 40
3.	Early White Spined 1 25	15	7. Man of Kent 3 00 30
			8. Small Gherkin (Burr) 4 00 4)

Kinds.—Nos. 1, 2, 3 and 4 are the earliest; No. 3 is the favorite kind for table use; Nos. 1 and 3 for pick ing, and 3 also for general use, very productive.

culture.—In February and March, plant seed in hot-bed; or, in April, in cold frame, well-covered. Plant out in rich, newly-dug ground, in hills or ridges, when night frosts are gone, and cover up with the St. John Hand Glass. In frames, protect well at night; give air in day; avoid winds; shade from mid-day sun. Keep up heat when grown early in the year by hot dung lining as temperature of bed subsides. Stop the leading shoots continually as the fruit shows; or plant the seed in open ground as soon as the weather becomes settled and warm, in hills four feet apart. Use a shovelful of warm manure to each hill, which cover with an inch or two of earth; scatter eight or ten seeds to a hill, cover half an inch deep with fine earth, and press it down. Hoe often, and when out of danger from insects thin the plants to four in hill.

CUCUMBER-ENGLISH VARIETIES FOR FRAMES.

₩ pkt.	
Giant of Armstadt 25	Cuthill's Black Spine 25
Lord Kenyon's Favorite 25	Eyre's Perpetual 25
Prize Fighter 25	Improved Sion House 25
'Walker's Rambler 25	White Bavarian 25
Telegraph 25	British Volunteer 25
Lynch's Star of the West 25	Invincible
Lancashire Witch 25	

DANDELION-PISSENLIT.

French Seed, 40 cents & ounce; \$4 \$ 10.

Culture.—The Dandelion is a hardy perennial plant, and one of the most desirable early Spring salads. Sow in May or June, in drills half an inch deep and ten inches apart; in July or August, thin out the plant to three inches apart. Keep clear of weeds during the Summer, and, in April of the ensuing Spring, the plants will be fit for the table.

EGG PLINT -MELONGENE - Berengena - Tierpflinge.

	7 15. 73 nz.		₩.	pkt.
1.	Long Purple \$4 00 40	D. St	riped (ornamental) \$0	10
2.	New York Purple 8 00 75	6. W	nite (prnamental)	10
3.	Black Pekin (fine) P pkt. 10	7. Ne	w Green from Thibet	10
4.	Scarlet Chinese (beautiful . 75 oz. 75			

Kinds.—No. 1 is the earliest, and very fine and productive. No. 2 is the targest and best for main crop; the others are used principally for ornament. No. 4 is most beautiful, and also a fine cooking variety. Nos 3 and 7 are new sorts, of excellent flavor and quite ornamental leaves.

Culture.—Sow in hot-bed, very early in Spring; transplant when two inches high into a second bed; if that is not done, thin to four inches apart. Do not plant out till the weather becomes perfectly settled and warm, and cover up with the St. John Hand Glass. Cool nights or wet weather will check them. Keep some back in frames for a second planting out, in case of weather changing unexpectedly. Keep plants watered for a few days if the sun is hot when put out. Keep plants tilled by the hoc, and draw earth up as for Cabbage.

ENDIVE—CHICOREE— Escarola ó Endivia—Endivien.

1. Green Curled \$3 00 30 30 Wild, or Chicory \$1 00 102. White Curled \$3 00 30 4. Broad-Leaved Escarolle) \$3 00 30

Kinds.—Nos. 1 and 2 are the best for Salad - particularly No. 1, which is also the most hardy. The roots of No. 3 are used for flavoring coffee.

culture.—Sow No. 4 early in Spring; Nos. 1 to 3, from latter part of May to end of July. Sow thin an loover slightly. When up, thin out to eight inches apart, and give a good watering afterwards, if dry. When leaves are six or eight inches long, blanch them by gathering the leaves in the land and tying them together near the top with yarn or bast. This must be done when quite dry or they will rot. At the approach of Winter the growing crop may be taken up carefully, with ball of earth to each plant, and placed close together in frame or cellar for use; but they must be kept dry, and have plenty of air, or they will rot.

GARLICK-AIL-Ajo-Rnoblauch.

Sets, 40 cents # 1b.

Culture. Light, rich soil. Plant the sets in rows six inches apart and about two inches deep. Leave a foot between the rows, which keep hoed deeply. When the leaves turn yellow, take up the crop.

ICE PLANT.

Pounce, 40 cents.

Culture. -- Sow in good garden soil as soon as ground becomes warm in Spring, in shallow drills—a desirable vegetable for boiling like Spinach, or for garnishing.

KALE—Chou vert frise—Col—Blättertohl.

Kinds.—Nos. 1, 2 and 5 are Borecole. No. 3 is a fine vegetable, having the appearance of white Celery, but cooked as Asparagus. No. 4 is used for early Spring Greens. No. 5 is a very tall and productive sort, highly recommended.

Culture.—Nos 1, 2, 5 and 6 are cultivated and grown exactly like Cabbage. No. 4 sow in September, for early Spring Greens. No. 3 sow early in Spring in a frame, transplant to another bed, six inches apart, in six weeks' time. Dig and manure highly a piece of ground, and plant out in hills three feet apart and three plants in each hill, about the end of July. Before Winter, cover these with litter or leaves. The following Spring uncover and dig in manure. Keep them well cultivated through the Summer, and the following Winter it will begin to produce a crop. Salt is a good manure to be given to this plant, as well as to Asparagus.

KOHLRABI, or Turnip-Rooted Cabbage—Chou RAVE—Colinabo.

- 3. Large White or Green " 2 00 " 20

Kinds.—Sow No. 1 for early forcing, and also for general crop. This is a delightful vegetable, and should be grown in every Garden.

Culture.—May be sown from April to July, and the plants afterwards treated the same as Cabbage, only, that in earthing up the plants, you must be careful not to cover up the globular part. Set out in drills one foot apart, and eight inches in the drill.

LEEK-POIREAU-Puerro-Borro.

1. Large American Flag	₽ oz.	40	cents.	P d.	\$4 00
2. American Giant Castle	44	40	46	44	4 00
3. Large Rouen	4.6	40	66	"	4 00

Culture.—Sow very early in Spring, in drills six inches apart and one inchedeep. Thin out to one inches apart. When they are about seven inches high plant them out in rows eight inches apart, and as deep as possible, so as not to cover the young centre leaves. Water thoroughly, if dry weather when planted out. Draw earth up to them as they grow. Require very rich soil. Take up and store away before Winter weather.

LETTUCE-LAITUE-Lechuga-Salat.

₩ lb. ₩	OZ.	智情 第02.
1. Early Wh. Forcing Head 34 00	40	11 Large Yellow Butter \$3 00 30
2. Early Curled Silesia 2 50	25	12. Brown Dutch 4 00 40
3. Curled Simpson 5 00	50	13. White Cos 4 00 4)
4. Large Perpignan 4 00	40	14. Green Cos 4 00 40
		15. Grey Cos 4 00 40
		16. Blood Cos 4 60 40
7. Speckled German 4 00	40	17. Oak-Leaved Cos 4 00 40
8. White Cabbage 3 00	30	18. Magnum Bonum Cos 4 00 40
9. Wheeler's Tom Thumb 5 00	50	19. Hardy Green Winter 3 00 30
10 Ice Drumheau 2 00	20	20. Brown Winter Cabbage. 4 00 40

Klads.—No. 1 is the best for forcing or earliest Spring crop. No. 3 for a succession kind. Nos. 5, 8, and 11—are five for main Spring and Soumer crops; No. 12 for Fall, and Nos. 19 and 20 for Win er. Nos. 13 and 14 are also highly recommended, but more trouble to raise.

Culture.—Sow Nos. 1, 2 and 3 in frames, in March, and in the open ground as early as it opens, and transplant in rows eight inches apart. Sow in a month's time same varieties again as also Cos for a succession. In August, sow any of the varieties, except Nos. 19 and 20, which should be sown in September. In October, some of these may be planted in frames to head in Winter and early Spring. Always sow thin, and thin out well, or the plants will not be strong. The last Spring sowing had better be grown where sown, being thinned out to six or eight inches apart. To have Cos in good order, they must be sown in hot-bed early in the year and transplanted in a cold frame, so as to have them good plants to set out at the opening of the ground. They require tying when grown, for a few days, to blanch. Lettuce requires good ground, and moisture in abundance. Hoe carefully every other week.

MARTYNIA-FOR PICKLES.

1. Craniolaria. Ptb. \$3 00; Poz. 30 | 2. Proboscidea. Ptb. \$3 00; Poz. 30 Kinds.—Both are productive and fine for pickles.

Culture.—Plant in May, in the open ground, three feet apart in each direction, where the plants are to remain, leaving only one plant in each hill; or the seed may be sown in hot-bed, and the seedlings afterwards transplanted.

MUSK MELON-MELON MUSCADE-Melon Muscatel-Kantalupen.

	\$ tb. \$`0	z.	 	Z.
L.	Fine White Japan \$4 00 4	0	8. Long Persian\$2 00 2	0
	Fine Nutmeg 1 25 1		9. Skillman's Netted 1 50 2	0
	New Valparaiso (fine) 4 00 4		10. Turk's Cap 1 50 2	0
	Orange Christina 3 00 3		11. Yellow Cantelope 1 50 2	
5.	Pine Apple 1 50 2	0.0	12. Pomegranate, for Perfume. 4 00 40	0
8.	Green Citron 1 25 1	.5		
7.	Jenny Lind 1 50 2	0		

ENGLISH VARIETIES FOR FRAMES.

Scarlet Gem	25	Camberwell Beauty Ppkt.	25
Orion "	25	Trentham Hybrid"	25

Kinds.—Nos. 1 and 7 are the earliest varieties. No. 1 is almost entirely skinless; small, but equal in flavor to the finest in cultivation. In addition to this sort for general crop, we recommend Nos. 2, 3 and 4.

Culture.—Plant in hills six feet apart each way, six seeds in each, and thin out to two or three plants when in a state of forwardness. It is a good plan to make a hole for each hill two feet wide, in which dig some rotten stablemanure or burnt sods. Dust a little soot or wood-ashes, recently burnt, on the growing plants, when up. Draw earth up around the stems, and stop their points to make them branch when they begin to run. Lay them out evenly to cover the ground, which keep clear from weeds at all times. A light but rich soil suits them best.

WATER-MELON-MELON D'EAU-Zandia-Baffermelonen.

	\$ 1b. \$ 02	P 1b. 8 02.
1.	Early Mountain Sprout\$2 00 20	6. Apple Seeded\$5 00 50
2.	Black Spanish 2 00 20	7. Large Green Virginia 2 00 20
3.	Ice Cream (fine) 1 25 15	8. Orange 4 00 40
4.	Goodwin's Imperial 4 00 40	9. Apple Pie (Citron variety)2 50 20
5.	Strawberry 4 00 40	10. Citron, for Preserves 2 00 20

Kinds,—No. 1 is early, and No. 3 is the finest for general crop. Nos. 5 and 7 are popular Southern sorts of excellent quality.

Culture. - Same as Musk-Melon, except that the hills should be eight feet apart each way.

MISHROOM-CHAMPIGNON - Seta-Champignonbrut.

English Spawn, 15 cents # tb.; 8 tbs for \$1 00.

Culture.—Mushroom-beds are be t made under cover. Any cellar or shed

will do. Collect a quantity of fresh stable-manure, without the long straw; turn it over daily for six or eight days to get rid of the rank heat. Dig out a foot and a half deep of the space to contain the bed. Lay some long manure at bottom, and then the prepared dung, a little at a time, evenly aud well beaten down, until it is a foot and a half high; put a layer of light earth on this, one inch thick; then another layer of dung, principally horse droppings, in eight or ten days, or as soon as the heat has declined to about 70°, or just to feel a little warm to the finger, insert lumps of Spawn the size of a hen's egg, ten inches apart, and just even with the surface of the bed. and cover with earth an inch thick. Beat it all gently down, and immediately inclose the bed all around, and over the top, with a wall of warm dung previously well prepared, by turning over to temper and sweeten it, and, if out of doors, keep off rain by mats or thin boards.

MUSTARD-MOUTARDE-Mostaza-Senf.

White London, 50 cents P quart; Black, 40 cents P quart.

Culture.—Sow thickly, during April and early in May, in shallow drills, and press the earth well down, and for Fall Salad, in September; or in frames or boxes, in any out-house, during winter. Sow very thickly.

NASTURTIUM-CAPUCINE-Maranuela-Nasturtium.

Tall, 20 cents \$\mathbb{H}\$ ounce, \$2 \$\mathbb{H}\$ lb.; Dwarf, 40 cents \$\mathbb{H}\$ ounce.

Culture.—Sow in May, or early in June, in drills about an inch deep; the Tall Kinds near fences or poles, on which they can climb and have support. If left to trail upon the ground the fruit is apt to be injured.

OKRA-Gombo-Quimbombo-Safran.

Improved Dwarf, \$\Psi\$ to., 75 cents; \$\Pounce\$, 10 cents. Long Green, \$\Pounce\$ to., 75 cents; \$\Pounce\$, 10 cents.

Cuiture.—Plant late in Spring, after the ground has become warm, in drills three feet apart, where the plants are to remain. Thin out to nine inches. Hoe, and draw the earth up occasionally to the stem. Should be well manured.

ONION - OIGNON - Cebolla - 3 wiebeln.

	ъ в в оz.	B bus. B qt.
1.	Early Red\$5 00 50	7. Red Top Onions \$7 00 30
2.	Large Red 4 00 40	8. Potato Onion 6 00 25
8.	Yellow Dutch 5 00 50	9. Yellow Onion Sets 10 00 40
4.	Yellow Danvers 5 00 50	10. White Onion Sets12 00 50
Б.	White Portugal 5 00 50	11. Shallots 7 00 25
6.	Welsh, (for Salad) 3 00 30	

Kinds.—Nos, 1 and 4 are the earliest; Nos. 2, 3, and 5 are the best for main crop; No 5 is also best for pickling, but should be sown very thick for this purpose.

Culture.—Rich soil, deeply dug, and then lightly rolled. Sow Nos. 1 to 5 moderately thick, in drills one inch deep and twelve inches apart. as soon as the ground is open in the Spring; thin to three inches apart. Keep down weeds, but avoid covering the bulbs. No. 6 should be sown last of August or early in September, and will produce new leaves for salad early in Spring. Onion Sets and Tops are placed on the surface, in shallow twelve inch drills, about four inches apart, but not covered. No. 8 plant in Spring or Fall, as they will survive the Winter if slightly covered with dung.

ORACH, or French Spinach.

Culture. - Sow early in Spring and cultivate like Spinach.

PARSLEY-PERSIL-Perijil-Beterfilie.

 1. Extra Curled
 # fb., \$1 50
 0 oz., 15

 2. Plain
 75
 10

Kinds.—No. 1 is the best; No. 3 the root only is used.

Cuiture.—Soak the seed a few hours in lukewarm water, and sow early in Spring, in drills an inch deep, and one foot apart. Thin out the plants to four inches apart. To preserve in winter, transplant some, and set them in a light cellar. No. 3 should be thinned out while young, and managed the same as Carrots and Parsnips.

PARSNIP-PANAIS-Chirivia-Baftinate.

1.	Sutton Student 1b	, 81	()()	₩ oz.,	15
2.	Guernsey, or Cup		75	66	10
3.	Early Round	1	50	66	20
4.	Long White "		75	56	10

Kinds.-Nos. 1 and 2 are the best-kinds. No. 3 is early, but small size.

Culture.—Sow as soon as ground opens in Spring, in drills one inch deep and fifteen inches apart. When plants are two or three inches high, thin out to seven inches in a row. Hoe often. Soil should be deep and rich.

PEAS-Pois-Chicaros 6 Guisantes-Erbien.

	EXTRA EARLY.		🕏 bus.	Pot.
*1.	Alpha (The New Extra Early Wrinkled Pea)	.3 feet.		
	in ½ pint packages, each \$3 00			
2.	Carter's "First Crop"	. 2 1 66	\$10 00	0 50
3.	Daniel O'Rourke	23 6	6.00	30
4.	Philadelphia Extra Early	21 .1	9 00	40
	EARLY		0 00	40
*5.	McLean's Advancer (fine)	.23 66	12 00	50
6.	Laxton's Prolific Long Pod (fine)	4 4		1 (0
7.	Warwick	.3 "	8 00	40
8.	Blue Surprise	4 66	10 00	50
9.	Rising Sun	3 11	10 00	50
10.	Laxton's Supreme (splendid Green Pea)		10 00	3 00
*11	McLean's Epicurean.	91 11		
14.	bichean's Epicutean,	, 49		1 00

Peas—(Continued.)	GENERAL CROP.	height.	₩ bus.	₩ qt.
13. Blue Imperial (fine). 14. Blue Scimetar. 15. Dwarf Sugar (edible 16. Tall Sugar (edible po *17. New Wrinkled Sugar *18. Hair's Dwarf Green 19. Waterloo Blue. 20. Harrison's Glory (fine 21. Harrison's Perfection *22. Napoleon (fine).	pods). ds). (edible pods). farrow. (fine).	. 2½ 6 . 3 "	\$09 00 6 00 8 00 14 00 16 00 20 00 10 00 8 00 10 00 8 00 10 00 10 00	\$0 40 30 50 60 60 80 50 40 50 40 50
	LATE CROP.			
 25. Hundred-Fold (New) *26. McLean's Wonderful 27. White Marrowfat 28. Black Eyed Marrowfa 29. Tall Matchless (fine) *30. British Queen *31. Epp's Monarch 	ne)		4 50 4 50 10 00 16 00 14 00 12 00	1 00 3 00 1 00 25 25 50 60 60
	VARIETIES-WITHOUT BRUSH.			
34. Tom Thumb, extra ear 35. Strawberry, early 36. Bishop's Dwarf, early 37. Bishop's Long Pod. ec 38. McLean's Princess Ro 39. The Peabody (fine br 40. Dwarf Waterloo (new 41. Queen of Dwarfs	(fine) extra early	.9 inches1 foot1 " .1½ " .1½ " .1½ " .1½ " .1½ "	14 00 10 00 9 00 9 00 9 00 9 00 6 50	50 50 50 50 50 40 1 00 1 00
42. Blue Eclipse		$.1\frac{1}{2}$ "	10 00	50

The above list comprises, it is believed, every desirable variety known, that has been tried, and found to succeed well in this climate.

Those marked * are Wrinkled Marrows, and should be planted thicker than the round Peas, as they are more liable to rot—they are the finest flavored of all the Peas.

Kinds.—We recommend, for the first crop, Nos. 2, 3, 4 and 33; for second crop, Nos. 5, 6, and 7; for general crop, Nos. 12, 13, 18, 22 and 23; for late crop, Nos. 28, 29, 30 and 32. For small gardens, Nos. 33, 34 and 41. No. 33 is one of the finest in cultivation.

culture.—A light, dry soil, not over rich, suits the Pea. If they grow too vigorously, and show no signs of bloom, run a spade down about six inches from the row, and thereby root-prune them. Do this each side the row, and they will bloom in a few days. Plant as early as the ground can be worked, and again every two weeks, for succession throughout the season. Plant in single or double rows, from four to six feet apart, according to the different heights, about an inch apart in the row, and three inches deep. In Summer, plant six or eight inches deep. Hoe often. In dry weather, Peas should be

soaked in soft water five or six hours before planting, and if the ground is very dry, they should be watered in the rows.

PEPPER -PIMENT -Pimiento - Pfeffer.

	P Ib. P	OZ.			P Ib. T	9 oz
1.	Red Cherry\$4 00	40	4.	Large Squash	.\$4 00	40
	Sweet Spanish for Salad) 5 00					
3.	Sweet Mountain (for Man-		6.	Long Cayenne (Yellow)	. 4 00	40
	goes) 5 00					

Kinds.-No. 1 is early; the others are best for main crop.

Calture.—Sow in hot-b-ds in March, or in open ground in a seed-bed in middle of Spring, in light, warm ground. When three mehes high, transplant to eighteen inches apart each way and cover with the St. John Hand Glass Hoe frequently.

POTATO -POMME DE TERRE--Patata--Rartoffel

The price may vary as the season advances. We will mail five pounds of any of these varieties, from No. 2 to 9 for \$1.

9 0	1000 000 000000, 31 0110 210 8 2 10 0, 301 42.		
		P Bus.	🛱 Barrel.
1.	"KING OF THE EARLY," or Brezee's No. 4. 7 lb., 75.	\$30 00	
2	EARLY ROSE, Blush skin and white flesh	. 2 50	5 00
3.	Early Goodrich, White skin and flesh	. 2 00	4 00
4.	Early Sovereign, Mealy when size of Walnuts	. 3 00	°7 00
5.	Early Cottage, White skin and flesh	. 3 00	6 00
6.	Early Dykeman, White skin and flesh	. 2 00	4 50
7.	Early Sebec, Smooth white skin and flesh	. 2 50	5 00
8.	Callao, Dark skin and pure white flesh	. 2 00	4 50
9.	Gleason, Blush skin, white flesh	. 2 00	4 50
10.	Calico, Mottled skin and white flesh	. 2 00	4 50
11.	Harrison, White skin and flesh	. 2 00	4 00

Kinds,—Nos. 1, 2, 3 and 4 are the earliest sorts; Nos. 2 and 3 are the most productive; Nos. 9, 10 and 11 are highly recommended for general crop.

Culture.—To grow healthy Potatoes and free from rot, it is necessary to plant them very early in the season, in good rich soil, that has been liberally manured the previous years. When the plants make their appearance above ground, apply to each hill a double-handful, each, of wood ashes and plaster mixed; if kept free from weeds, a large crop will be the result. To have Potatoes VERY EARLY, the seed must be sproute l, i.e., it must have made an advance. ment in growth of at least an inch. This sprout should be strong and well developed, its thickness about that of the steam of a common tobacco pipe, and its top crowned with green buds just bursting into leaf; at the bottom of the sprout are emitted, or in the course of emission, small, thread-like roots, which, as soon as planted, take possession of the soil. Here is the great secret of obtaining early Potatoes.

To acquire this stage of advancement, many expedients are resorted to. It is the natural habit to sprout early; indeed, oftentimes there is a difficulty in

retarding this movement, for if the sprout is too long, it runs great risk of being knocked off in course of setting. The mode generally adopted is, in January and February, place them in baskets, not over a peck in each, in a very warm garret, or other room, that is light and warm.

PUMPKIN-CITROUILLE-Calabaza-Croße Rürbis

% lb. % oz.	Ph. Poz
1. Large Cheese\$0 60 10	3. Mammoth\$2 00 20
2. Cushaw 1 00 10	4. Connecticut Field # qt. 40
KindsNo. 1 is the best for family us	se; No. 4 for stock.

Culture.—May be planted middle of Spring, amongst the Indian Corn, or in the field or garden, in hills eight or ten feet apart each way, four seeds in a hill. In other respects, are cultivated in the same manner as Melons and Cucumbers; but avoid planting them anywhere near other vines.

RADISM--Radis et Rave--Rabanos and Rabanitos--Nadies und Rettig.

₩ 15. ₩ oz.	₩ 1b. ₩ 0z.
1. French Breakfast \$1 50 15	13. White Summer Turnip\$1 25 15
2. Early Frame 1 00 10	14. Yellow do do 1 50 15
3. Early Scarlet Turnip 1 00 10	15. Purple do do 1 50 15
4. Early White Turnip 1 50 15	16. Grey do do 1 50 15
5. Early Yellow Turnip 1 50 15	17. Black Spanish 1 25 15
6. Rose, Olive-Shaped 1 00 10	18. White Spanish 1 50 15
7. Scarlet do 1 00 10	19. Scarlet Chinese Winter. 2 00 20
8. White do 2 00 20	20. Purple do do 30
9. Long Scarlet Short Top. 75 10	21. White do do 30
10. Long Salmon 1 00 10	22. Raphanus caudatus, or
11. Long Purple 1 50 15	edible pod Radish P pkt. 10
12. Long White Naples 1 50 15	

Kinds.—Nos, 1, 2, 3 and 9 are best for early sowing; Nos. 3, 7 and 9 for succession; Nos. 12, 13, 14 and 16 are best for Summer sowing; Nos. 17, 18 and 19 for Winter use; No. 19 is the best sort—keeps like a Turnip.

culture.—Sow Nos. from 1 to 10 as early in Spring as ground can be worked and every two weeks till the 15th of June, from which time sow Nos. 12 to 16 till 1st of September. Nos. from 17 to 21 should be sown from 1st of August to to the 1st of September. A light, rich soil suits best.

RHUBARB—RHUBERBE—Ruibarbo—Rhabarber.

Roots, 25c. to 50c. cach; \$3 to \$5 \$\pi dozen. | Linnæus, 30c. \$\pi oz; \$3 \$\pi tb.

Kinds.—We have found the above sort so much superior to all other in quality, size and earliness, that we have cultivated only this one sort.

Culture.—Sow in drills an inch deep; thin out to six inches apart; in the Fall trench a piece of ground, and manure it well; then transplant the young

plants into it, three feet apart each way; cover with leaves or litter the first Winter, and a dressing of coarse manure should be given every Fall; through the Summer, when the weather is dry, water once a week with soap-suds.

ROQUETTE.

30 cents \$\mathfrak{P}\ ounce.

Culture. Sow early in Spring in rows, and thin out to eight or ten inches.

SALSIFY, or Vegetable Oyster-Salsifis-Ostion Vegetal—Haferwurzel. Long White, P. d., \$4 00; P. oz., 40 cents.

Culture.—Sow as early as the ground can be worked in the Spring, in drills twelve inches apart. one inch deep, and thin out to six inches in a row. Keep them clean from weeds. Cultivate the same as for Carrots and Parsnips.

SAVOY.

Feather Stem Savoy, 40 cents per ounce.

A new Hybrid, possessing the habits of Brussels Sprouts, a delicate and delicious vegetable; requires the same treatment as Brussels Sprouts.

SCOLIME.

An entirely new vegetable from Spain. The root resembles a small Parsnip, and will keep through the Winter, like a Turnip. It is cultivated precisely like the Carrot, and boiled like Salsify. Seed expected. 25 cents packet.

SCORZONERA—Scorzonere—Escorconera—Scorzionerwurzel.

30 cents Pounce; \$3 00 P pound.

Culture.—Cultivate precisely in the same manner as Salsify.

SKIRRET—CHERVIS—Buderwurzel.

30 cents # ounce.

Culture.—Sow in drills, half an inch deep and a foot apart; thin out to eight inches in a row; keep down weeds. Take up the roots before Winter, and store in sand or earth.

SORREL-Osseille-Acedera-Sauerampfer.

Garden, 15 cents & jounce; \$1 50 p pound fulture.—Sow in Spring, in drills six or eight inches apart.

SPINACH-EPINARDS-Espinaca-Epinat.

	₹ 1b. €			. \$ 02.
1. Flandre (fine)	.\$0 60	10	4. Yellow Sorrel-Leaved \$0	60 10
2. Round-Leaved	. 60	10	5. Prickly	60 10
3. New Zealand (Summer)). 2 00	20		75 10

Kinds,—Nos. 1 and 2 are the best for early Spring sowing; No. 2 also for

Fall and Winter, in this latitude. No. 5 is the most hardy.

Culture.—Rich soil; if light press or roll it down before sowing. Sow a quarter of an inch deep, and a foot between the rows. Thin out to 8 inches. Sow as soon as frost is out of ground, and hoe frequently to prevent the plants running to seed. For Winter Crop, sowend of August and in September. Lay straw between the rows at approach of Winter, and very thinly over the leaves.

SQUASH-GIRAUMON-Calabaza-Rüchenfürbis.

1. Early Golden Bush	8. Hubbard (fine). \$2 50 30 9. Winter Crook-Neck. 1 25 10 10. Turbon. 3 00 30
5. Yokohama (fine). 3 00 30 6. Lima Cocoa-Nut. 2 00 20 7. Boston Marrow 2 00 20	11. Algora

Kinds.—Nos from 1 to 4 are the best sorts for early Summer use; Nos. from 5 to 10 are the best for late and Winter use; Nos. 5, 7 and 8 are particularly recommended; Nos. 11 and 12 are fancy sorts.

Culture.—Plant in hills, in the same manner, and at the same time as Cucumbers and Melons; the Rush three or four feet apart, and the running kinds, from six to nine.

TOBACCO SEED.

Ph. Poz.	P 16. B 0z.
Imported Havana\$10 00 75	Ruffled-Leaved Virginia \$6 00 50
Connecticut Seed Leaf 4 00 40	James River Virginia 6 00 50
Maryland 5 00 40	
Florida 6 00 50	Kentucky 6 00 50
Ohio 6 00 50	

TOMATO TOMATE Tomates - Liebesapfel.

	P Ib. F	oz.	₩ 1b. ₩ 0z
1.	Early Red\$3 00	30	9. Yellow Pium\$5 00 50
2.	Large Red Smooth 3 00	30	10. Red Plum 5 00 50
3.	Lester's Perfected 4 00	40	11. Large Yellow 4 00 40
4.	Fejee Island 4 00	40	12. Red Cherry 5 00 50
5.	Cook's Favorite 4 00	40	13. Yellow Cherry 5 00 50
6.	Tilden 4 00	40	14. Pear Shaped 5 00 50
7.	Key's Prolific 3 00	30	15. Tomate de Laye # pkt., 10
	Cedar Hill 5 00		16. Strawberry 16., 5 00 50

Minds.—No. 1 is the earliest; Nos. 2, 3, 5 and 6 best for main crop; Nos. 9, 10, 11, 12 and 13 for pickles and preserves; Nos. 4 and 7 are valuable sorts for general use; very solid. No. 15 is the French upright variety; entirely distinct in its habit of growth, and of fine quality. No. 16 is the Winter Cherry sometimes called Strawberry Tomato; it is excellent for preserves.

Callure.—Sow in hot-bed in March, and transplant into the open ground when weather becomes warm and settled; or sow into open ground in May, and transplant when plants are six inches high. They succeed best near a fence or trellis to climb on, but they do very well if planted four feet distant from each other without support. Soil should be made rich and light.

TURNIP—Navet—Nabo—Rüben.							
% Њ. № oz. % Нъ. № оз							
1.	Early Dutch	0 75	10	12.	Yellow French (fine)\$0	75	10
	German Teltow		20	13.	Yellow Stone	75	10
3.	Cow Horn	75	10	14.	Yellow Aberdeen	75	10
4.	Red Top Strap-Leaf	75	10	15.	Robson's Golden Ball	75	10
	White Norfolk	75	10	16.	Yellow Finland 2	00	20
	White Strap-Leaf Flat	75	10	17.	Dale's Hybrid	75	10
7.	White Flat or Globe	75	10	18.	Improved Yellow Sweed-		
8.	White French	75	10		ish, or Ruta Baga,	75	10
9.	Green Globe	75	10	19.	White Ruta Baga	75	10
10.	Long White Tankard	75	10		Laing's Ruta Baga	75	10
	Vertue's Long White	1 00	10	21.	Skirving's Ruta Baga	75	10
	0				ng sowing; > os. 3, 4, 6, 8	3. 12,	14
					15 are also fine for table u		

Cuitare .- For early Tur ips. sow as soon as ground opens in Spring. Nos. 8, 18, 10 and 21 should be sown from any time in July to first of August; but the other kinds, for Winter use, may be sown from middle of July to end of August. Turnips are generally sown broadcast, but much larger crops are obtained (particularly of the Ruta Raga) by cultivating them in drills twelve inches apart, and the ned to 6 nches in the drills. Hoe them a month after sowing.

VARIOUS KINDS OF VEGETABLES USED FOR PICKLES. DIRECTIONS FOR CULTURE OF EACH WILL BE FOUND UNDER THE PROPER HEAD.

Refugee Beans	# bus.	\$6 00	# qt. \$	30
Blood Turnip Beet	7 lb.	1 00	Toz.	0 15
Brussels Sprouts	6.6	3 00	66	0 30
Red Dutch Cabbage	66	4 00	6.6	0 40
Early Paris Cauliflower	6.6	15 00	66	1 50
White Spined Cucumber		1 25	44	0 15
Gherkin	66	4 00	66	0 40
Martynia	66	3 00	4.6	0 30
Mango Melon	6.6	2 00	61	0 20
Scarlet Gem Melon			P pkt.	0 25
Tall Nasturtium	# Ib.	\$2 00	Poz.	0 20
White Portugal Onion		5 00		0 50
Sweet Mountain Pepper		5 00	6.6	0 50
Cherry do		4 00	6.6	0 40
Strawberry Tomato		5 00	4 6	0 50
Red Cherry do		5 00	4 4	0 50
Yellow do. do		5 00	6.6	0 50

VARIOUS KINDS OF VEGETABLES USED FOR SALADS.

- 1	Burnet			9	OZ.	20	Endive	5 Ib.	\$3	00	P oz.	30
3	Borage	# Ib.	\$.	50	4.6	25	Lettuce	46	3	00	16	30
	Celery						Mustard				뭐 qt.	50
4	Corn Salad	66					Swt. Spanish Pepper		5	00	Poz.	50
-	Cress	6.6		60	44	10	Roquette	66	3	00	66	30
1	Dandelion	66	4	00	6	40	Sorral	66	- 1	5.0	66	15

TREE AND SHRUB SEEDS.

For Description, and many other Kinds, see our Tree and Shrub Seed Catalogue In quantities not less than quoted.

and quantities for the train quantities				
Abies, Alba, White Spruce	₩ lb.	\$5 00	W oz.	50
" balsamea, Balsam Fir	"	3 00	66	30
" Canadensis or Hemlock Spruce (clean seed)	66	5 00	66	50
" excelsa, or Norway Spruce	66	1 00	66	20
" Fraseri, or Double Spruce	66	6 00	4.6	60
" pectinata, or European Silver Fir	6.6	1 25	66	20
" Nordmanniana, or Nordman's Spruce	66	10 00	66	75
Acer plantanoides, or Norway Maple		1 50	46	20
" pseudo-platanus, or Sycamore Maple	44	1 50	66	
pocado pidaldas, or egoanoro acaptorititi	6.6	2 00	44	20
Allanda same wis an Alder	66		66	25
Alnus communis, or Alder	66	1 00	66	20
Berberis vulgaris, or Common Barbery	66	2 00		25
Betula populifolia, or Poplar-Leaved Birch	66	.2 50	"	30
" alba. or White Birch	66	2 00	44	20
Buxus sempervirens or Evergreen Box		5 00	6.6	50
Catalpa bignonioides, or Catalpa	66	6 00	66	60
Cedrus Libani. or Cedar of Lebanon			cone,	30
Cornus paniculata, or Panicled Dogwood	66		Poz.	30
Cupressus sempervirens, or Common Cypress	66	3 00	44	30
" disticha, or Diciduous Cypress	66	1 00	66	20
" horizontalis or Horizontal	66	4 00	66	40
Diospyrus Virginiana, or Persimmon	66	2 50	66	30
Fraxinus alba, or White Ash	66	1 50	66	15
" ornus, or Flowering Ash	46	3 00	6.6	30
Ginkgo biloba. or Salisburia.		₩ doz.		75
Gleditschia, triacanthos, or Honey Locust	66			
Gymnocladus Canadensis. or Kentucky Coffee Tree	66	3 00	₩ oz.	10 30
	- 66	50	66	
Juniperus prostrata, or Prostrate Cedar	"		6.6	10
" communis. or English Juniper	44	1 50	"	20
Larix Europea, or European Larch	"	2 00	••	25
Maclura aurantiaca, Osage Orange, ₩ bus. 33 fbs., \$18	66	75		
Magnolia acuminata, or Cucamber Magnolia		3 00	66	30
" grandiflora, or Large Flowering Magnolia	44	4 00	44	40°
Morus alba, or White Mulberry	6.6	3 00	66	30
Paulownia Imperialis, or Paulownia	66	5 00	4.6	50
Pinus maritima, or Sea-Side Pine	4.6	1 50	6.6	15
" alepica or Swiss Pine	66	4 00	44	40
" laricia, or Corsican Pine	66	4 00	44	40
" nigra Austriaca, or Black Austrian Pine	66	2 00	6.6	25
" rigida, or Pi/ch Pine	6.6	5 00	66	50
" Pinea. or Stone Pine	44	2 00	4.6	25
" cembro. or Cembran Pine	6.6	1 25	64	20
" Lambertiana, from California	66	12 00	" 1	00
" Sabiniana, or Sabine's from California	66	18 00	" 1	50
strobus. or Weymouth Pine	66	4 00		40
" sylvestris or Scotch For	66	2 50	66	
5,1.050115, 01 2.00000 2.7	6.6	1 00	44	25
Rhamnus catharticus, or Buckthorn	64		46	15
Rhus cotinus, or Purple Fringe	66	6 00	"	60
Robinia pseudoacacia or Yellow Locust	"	1 00		15
Taxus bacata, or English Yew		4 00	"	40
Thuja occidentalis, or American Arbor Vitæ (clean seed)	6.6	4 00	44	50
" orientalis, or Chinese do	. 66	4 00	"	40
Tilia Americana or American Linden	44	1 00	4.6	10
Ulmus Americana, or American_Elm	"	3 00	6.6	30
Zanthoxylum Americanum, or Prickly Ash	66	5 00	6.6	50

Pot, Sweet, and Medicinal Herb Seeds.

Charles and TVT-street Name of Street	le le Cusine en Oferiferentes
Graines d Herbes a l'usage d	le la Cusine ou Odoriferentes.
Dill	Anethum graveolens 9 oz. 15
Sweet Fennel	TOUR THE TOUR TOUR TOUR TOUR TOUR TOUR TOUR TOUR
Garden Angelica	Archangelica officinalis " 15
Arnica	Artamicia abainthium
Wormwood	Artemisia absintinum
Belladonna	Atropa Belladona " 75
Borage	Borago officinalis " 25
Pot Marigold	Calendula officinalis " 30
Saffron	Carthamus tinctorius " 15
Hemlock	Conium maculatum " 50
Carraway	Carum Carvi " 15
Coriander	Coriandrum sativum " 15
Purple Foxglove	Digitalis purpurea " 40
Boneset	Eupatorium perfoliatum " 80
Liquorice	Glycyrrhiza glabra " 75
Pennyroyal	Hedeoma pulegioides " 75
Henbane	1 0
Elecampane	
Hyssop	
Lavender	Lavendula vera 40
Dandelion	Leontodon taraxacum 40
Horehound	mailubium vuigale 1 00
Balm	Inclissa omornans
Catnep	Nepata catama
Sweet Basil	Ocymum basilicum " 50
Dwarf or Bush'd Busil	minimum · 60
Speet Marioram	Origanum mariorana " 50
Pot Marjoram	" onites " 75
Pot Marjoram	Ricinus communis
Rosemary	Rosamarinus \$ oz. 50
Dyer's Madder	
Rue	
Red Sage	Salvia hominum. " 50
Common Sign	" offic nalis " 30
Common Sage	" offic nalis " 30 Satureja hortensis " 30
Pone	Sesamum orientale
Bene.	cesamam orientate
Tansy	
Thyme	Thymus vulgaris
Broan-Leavea English Thyme	" 60
Anise	Trajium anisum
Fenugreek	
Pear Seed P b. \$4 P oz \$0 40	SEEDS.
Pear Seed 7 fb. \$1 7 oz \$0 40	Red Alpine Strawberry Foz. \$1 50
Apple Seed & bus \$12.7 qt. 0 50	Strawb'y Seed 10 varieties " 2 00
	Black Mazzard Cherry Pits 79 at. 0 50
Currant Seed 0 50	Mahaleb Cherry Pits " 0 60
	Apricot Pits " 0 75
Goos berry Seed 1 50	Nectarine Pits 7 fb. 1 50
Mixed Strawberry ' 1 50	Peach Pits 3 bus. \$2 00. 7 qt. 0 10
EOD H	EDCES
Osage Orange	Maclura aurantiaca 7 lb. \$0 75
Hawthorn.	Cratægus oxycantha " 1 50
Ho ey Locust, or Three-thorned Acacia.	Gleditschia triacanthos " 75
Buckthorn	
Yellow Locust	Robinia pseudo-acacia " 1 00
Prince Tanan	Liquetrum Japaniaum 66 3 00
Privet Japan	Lighthum Japonicum 3 00
Sheep Berry	Viburnum prunifolium " 2 00
American Arbor Vitæ. Barbery	Thuja Occidentalis 4 00
Darvery	Berberis vulgaris

~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~
GRASS SEI	EDS.
Agrostis stolonifera, or Creeping Bent Grass a v	valuable lawn variety (nothing
better for grass-plots.).  vulgaris, or Red-Top Grass, \$\mathcal{B}\$ sack, \$5  Aira cæspilosa, or Hassock Grass. This Grass p	\$5.00 \ bus.
vulgaris, or Red-Top Grass, \$\mathbb{B}\text{ sack, \$5}	\$1.50 "
Aira cæspitosa, or Hassock Grass. This Grass p	orefers stiff or marshy bottoms,
where the water stands	
flexuosa, or Wood-Hair Grass,-delights on h	igh and rocky hills, 50 cts. 39. th.
Alopecurus pratensis, or Meadow Foxtail-one of	the hest Grasses for perma-
nent pasture, and thrives in perfection	on lands of an intermediate
quality as to moisture and dryness	60 cents % th
agrestis or Slander Foytail	75 cents 9 th
Anthoxanthum odoratum, or Sweet-Scented Verna	Cross 75 cents (1) th
Arundo avenaria, or Reed Grass	Of conta the
Avena elation or Tall Manday Oct Cross	2450 CO has
Avena elatior, or Tall-Meadow Oat Grass.  flavescens, or Yellow-Oat Grass. This G	man should be some with other
Juvescens, or remov-out Grass. This G	rass should be sown with other
varieties, such as Crested Dogstail and S	weet vernal-valuable for dry
meadows and pastures	\$1.25 😝 10.
Bromus Schraderi, or Rescue Grass. Cynosurus cristatus, or Crested Dogstail—a des	
Cynosurus cristatus, or Crested Dogstail—a des	arable Sheep-Grass and very
productive.  Dactylis glomerata, or Orchard Grass.  Festuca duriuscula, or Hard Fescue—one of the	
Dactylis glomerata, or Orchard Grass	\$250 \$\ bus.
Festuca duriuscula, or Hard Fescue—one of the	finest of the dwarf-growing
Grasses, and succeeds best on light, rich	soils35 cents # ID.
loliacea, or Darnel-spiked Fescue. This	Frass possesses all the valuable
properties of Rye Grass; its produce is	s larger, it springs earlier, and
improves by age, which is not so with Cor	mmon Rye Grass. 70 cents # fb.
ovina, or Sheep's Fescue-a good variety	for forming grass-plots; it has
the advantage of a fine and succulent foli	age35 cents \$ 1b.
pratensis, or Meadow Fescue. This Gra	ass makes excellent hay, and,
though a large plant, yet the herbage is s	succulent and tender, and much
relished by cattle, as it does not for	m rank tufts like the longer
Grasses	60 cents # 1b.
Grasses.	s much affinity to the Festuca
duriuscula,-from which, however, it is	distinguished by the leaves.
which are broader and longer.	\$1.00 % fb
which are broader and longertenuifolia, or Slender Fescue—flourishes	well on dry and sterile soils
to the state of th	[50 cents # lb.
Holcus lanatus, or Meadow Soft Grass-grows f	
thrive in any soil, from the richest to the	
Lolium Italicum, or Italian Rye Grass	
perenne, or English Rye Grass	\$4.00 ''
Onobruchis sativa or Saintfoin	\$5.00 "
Onobrychis sativa, or Saintfoin.  Panicuim Germanicum, or Hungarian Grass  Phalars arundinacea, or Reed Canary Grassg	\$3.50 "
Phalars aryndinassa or Road Capary Grass or	rows wall by the side of rivers
and standing pools	75 cents 30 th
Phleum gratence or Timothy	Market price
Phleum pratense, or Timothy	an excellent posture wariety
for work wat situations	75 cents 40 th
for very wet situations	a sorliest and awastest parture
annua, or Annual Spear Grass—one of the	earnest and sweetest pasture
varietiesnemoralis, or Wood-Meadow Grass. The	a combine mounth of this Crows in
nemoralis, or wood-Meadow Grass. The	early growth of this Grass in
the Spring, and its remarkable fine such	scurent and nutrative nerbage,
recommend it strongly for pastures	(14 Har)
recommend it strongly for pastures  pratensis, or Kentucky Blue-Grass  do. do. extra	(14 108.) 55.00 & Dus.
do. do. do. extra	stean(14 los.)\$5.50
trivialis, or Rough-Stalked Meadow Gra	ass. The superior product of
this Grass over many other species, its	nighty nutrative qualities, and
the marked partiality which oxen, horse	
merits which distinguish it as one of the	most valuable of those Grasses,
which affect moist, rich soils and shelter	
Mixed Lawn Grass (sow three bushels to the acr	(e)\$350 \ bus.
French Mixed Lawn Grass (sow three bushels t	o the acre)\$5.00 "

#### Formation of Lawns from Seed.

In the first place, see that your ground is thoroughly drained and well prepared, sowing at least three bushels of seed to the acre.

Perhaps the best description of soil for a lawn is the stiff loam or clayey soils which predominate in so many districts. This ought not to be by any means too rich, as a rapid growth is not wanted in the grasses of a lawn; but, in preparing it for laying down, let the surface be as much alike in quality as possible, and do not stint the quantity of seed. A very stiff clay is no better than a very dry sand for resisting drought, as it is, in a measure, sealed up against the insertion of roots, and the surface contracting by the withdrawal of moisture, it is liable to crack, &c., to a great depth. Nevertheless, a stiff soil usually makes the best lawn.

If your ground is in condition to raise a good crop of Potatoes, it will make a good lawn, adding about half a ton of Bone Dust to the acre, harrowed in before sowing the seed. The success of establishing a good lawn depends much on the season and condition of the ground at the time of sowing. A very slight raking in will protect the seed much, or a sowing of wood ashes will render them distasteful to birds. Rolling however, is at all times indispensable, and it is a good practice to sow a very thin scattering of oats amongst the grass seeds, which, coming up quickly, tends to shade and protect the tender grass.

The best time to sow grass seed is early in the Spring or about the beginning of September. If at the latter time, the seeds ought to have been the produce of the same season. Sowing plenty of them is also advised as the little extra expense for a good lawn ought not to be denied.

One important thing should not be forgotten in the preparation of the ground: let it all have a surface of about six inches alike, for nothing looks worse than to see a lawn grow all in patches. One exception however may be mentioned, and that is, if there be any steep slopes facing the South, or other exposed places, let the earth on them be better than in the ground level, for the aspect and other causes render such places, liable to burn with less sun than level places. They ought, therefore, to be of a better material, and sod ought to be provided to lay there if accuracy be expected. When the grass becomes well established, it should be frequently cut; the oftener you mow, the finer your grass will be.

It is only proper here to observe, that no lawn can be maintained long in good order without successive rollings. Mowing alone will not secure a good bottom without that compression which the roller tends to give.

## Graines pour les Oiseaux.

Prices variable—by the Quart or Bushel.			
CanaryPhalaris canariensisal	bout 3	9 qt. 2	5
HempCannabis sativa	66	" 2	0
MawPapayer rheas	66 Q	9 Hb. 3	0
Millet Panicum miliaceum.	·	g at. 2	0
RapeBrassica napus	66	" 3	0
Rough Rice Oryza sativa	44	66 3	0

#### Lilium Auratum,

\$5,00 \$ doz. 50 cents each.

## CLOVERS.

Yellow Trefoil	Medicago Lupulina	₩ lb.	60
Lucerne, or French Clover			50
Red Clover			20
Large Red or Pea-Vine		6 6	25
White Clover	repens	46	65
Scarlet Clover	incarnatum	44	60
Alsike Clover	hybridum	66	75
Bokhara Clover		66	75

## ESCULENT ROOTS AND PLANTS.

Plantes et Racines Potageres.

Asparagus Roots, Giant\$1 00 \$\mathbb{R}\$ 100\$8 00 \$\mathbb{R}\$ 1,000
Do. do. Conover's Colossal\$3 00 \$\mathref{100}\$ 100. \$25 00 \$\mathref{1}\$ 1 000
Horse Radish Roots
Rhubarb Roots
Dioscorea Batata, or Chinese Yam
Jerusalem Artichoke
Tarragon Plants
Earth Plants

CABBAGE, CAULIFLOWER, BROCOLI, BRUSSELS SPROUTS, CELERY, TOMATO, EGG PLANTS. Etc., E.c., at the proper Season.

## MISCELLANEOUS.

Ramie Seed # 1b. \$25 # oz. 2 0	0   Opium Poppy 滑 tb., 2 00
Fine English Oats B bus. \$3 0	
Caper Tree	
Chicory Seed	0   Cuba Bast (for tying) " 75
Chinese Sugar Cane # qt. 3	0 Japan Fibre (for tying) ' 1 25
Broom-Corn	
Sun-Flower 5	
Sea-Island Cotton # 1b. 3	0   Spurry # lb. 40
Tennessee Upland Cotton " 2	
Flax Seed	SpringVetches, \$\mathfrak{H}\ \text{bus.\$4.50} \mathfrak{H}\ \text{qt.}  25
Madder	Indelible Garden Pencils ea. 25 to 1.25
Seradella	Round Scotch Scythe-stones, each 30

Archangel Mats, \$1 each; \$9 7 doz.

## COLLECTIONS OF FLOWER SEEDS.

100	TT 1	0 4 1 7		7.70	1 0		
100	Varieties	of Annuals, E	gienniels	and Perennia	als, for	,	. \$5.00
50	do.	do.	do.	do.	66		. 2.50
20	do.	Annuals.					
10	do.	do.			68		. 50
20	do.	More Rare	Annuals,		6.6		2.00
10	do.	do.	do.		66		. 1.00
50	do.	Choice Gre	en-House	Seeds	6.6		8.00
25	do.	do.	do.	do.	166		. 4.00
20	do.	Hardy Bier	nials an	d Perennials			. 1.00
	~	ATT STRICT	TT TO OT	D OTHER PET	TOME	O PTG -DG	

BY ALL STRICTLY OUR OWN SELECTIONS.

## Day's Indelible Garden Pencil,

In India Rubber Cases; always ready; require no sharpening; each Pencil has an extra bex of 4 leads. Price, \$1 25 each.

# NOVELTIES, &c.

	ts. 50
French Yard Long Beans, entirely distinct from the Asparagus or Yard Long Beans; we consider this one of the very best Pole Beans in cultivation, \$\partial qt \qquad \text{this one of the very best Pole Beans in } \\$1	00
Marshall's New Yellow Pole Beans, very delicate and good flavored, \$\partial qt. \qquad \tag{\$1}	00
Blue Podded Runner Beans. This valuable variety, introduced last year, produces purple Flowers, and purple Pods, very ornamental, and also one of the best table varieties, #qt\$1	00
New Egyptian Blood Turnip Beet, small, very smooth, and handsome shaped dark Beet, of excellent quality for the table, \$\Pi\$ oz	50
New Dwarf Brussels Sprouts, growing a third less in height than the old variety, and superior flavor, $\textcircled{B}$ oz	50
Marblehead Mammoth Cabbage. Although not a new variety, we would call the attention of all wanting extra large heads to this sort,   → oz.,	75
*Crosby's Early Sweet Corn. Said to be the best Early Sugar Corn now in cultivation, Pqt	50
New Valparaiso Musk Melon. A new dark green and very large Melon of excellent quality, 😜 oz	40
Alpha Peas. A new "Extra Early" blue wrinkled variety of exquisite flavor, said to be as early as the earliest, and very productive. In ½ pint puckages, each. \$3	00
Hundred Fold Pea. A new first-class sort for main crop, productive and beautiful olive-green round Pea, ₽ qt\$3	00
Queen of Dwarf Pea. Not a new kind, but one that should be more generally cultivated. This Pea is of a branching habit, and grows but 1½ foot high, very strong haulm and requiring no support, ₽ qt.	50
King of the Early Potato. Claimed to be a week earlier than the "Early Rose," and for which \$50 apiece was asked last season.	75
Scolime. The attention of Horticulturists has recently been called to this interesting edible rooted plant from Spain. The mode of culture is the same as for Carrots and Parsnips, but should not be sown in this latitude until the first of June. The roots will keep during Winter, like a Turnip; should be cooked and served like Salsify.	25
Mammoth Chili Squash. Very large and interesting—seeds from specimens weighing over 200 lbs. each, #P packet of 10 seeds	25
Currant Tomato. The most ornamental of all the sorts. It is a veritable  Tomato. with long racemes of fruit of a bright red color, and not larger than the "Cherry Currant," very interesting and ornamental,  packet.	25

······	~
HORTICULTURAL IMPLEMENTS.	
AVARANCATORS, OR POLE PRUNING SHEARS \$3	00
VINE SCISSORS - For thinning out Grapes	50
FLOWER GATHERERS.— Scissors combining Tweezers and	
Pincers 1 50 to 2	50
	00
	00
	50
	00
CITIES DOLLE LIVE TO THE CONTROL DOLLERS CONTROL DOLLARS CONTROL CONTROL DOLLARS CONTROL CONTR	50
	50
	00
	00
	25
TRANSPLANTING TROWELS, - (Best cast-steel.) 1 00 to 2	00
DO. DO. (Black) 40 to	60
DUTCH, or PUSH HOES—(Cast-steel) from 2 to 10 inches 40 to 2	00
DO. DO. (German Steel), from 3 to 10 in. 35 to 1	50
TRIANGULAR HOES.—Are also used as Tree Scrapers 50 to	75
LADIES' GARDEN HOES	75 75
CORN HOES	50
LAWN RAKES.—Are used to cut off the flower-heads or buds	00
	00
	0.0
	00
GREEN-HOUSE SYRINGES.—(Brass), Pool's best make 4 00 to 11	00
GRASS HOOKS or SICKLES - Three sizes. 75 to 1	50
	00
BRUSH HOOKS	00
BRUSH SCYTHES. 2	00
AMES' C. S. BRIGHT SPADES. 1 DO. DO. DO. Long-handled. 1	75 75
	25
AMES' BEST C. S. BRIGHT SHOVELS	75
DO. DO. DO. Pointed long handle.  BEST SPADING FORKS	75
BEST SPADING FORKS.	25
ENGLISH PATENT C. S. LAWN-SCYTHES	75
SAYNOR & COOKE'S CELEBRATED PRUNING KNIVES, 2 00 to 3	00
SAYNOR & COOKE'S CELEBRATED BUDDING KNIVES, 2 00 to 2	50
FRUIT GATHERERS	40
ROUND SCOTCH SCYTHE STONES	30
BOOKS.	
Gardening for the South (Wm. N. White) \$2	00
The Young Gardener's Assistant(Thomas Bridgeman) 2	50
The Kitchen Gardener's Instructor. (Thomas Bridgeman)  Breck's Book of Flowers. (Breck) 1	75
	75 50
Grasses and Forage Plants(C. L. Flint) 2	50
	25
Fuller's Grape Culture	50
Florer Garden Directory (P. Print) 1	50
Gardening for Profit(P. Henderson) 1	50
Forest Tree Culturist (Fuller) 1	50
Practical Floriculture(P. Henderson) 1	50
Small Fruit Culturist	50
Miniature Fruit Garden(Thomas Rivers) 1	00

# Complete Assortments of Kitchen Carden Seeds for Private Families.

Choice Sorts	Most Approved and in Quantities to suit arge Gardens.	No. 1. \$5.	No. 2. \$10.	No. 3. \$18.	No. 4. \$25.
	Death Circ Wooden	1 pt.	1 qt.	1 qt.	D. or b
Beans	Early Six Weeks	1 pt.	I qu.	1 qt.	2 qt. 1 qt.
	Refugee	½ pt.	1 pt.	1 qt.	1 qt.
*	Horticultural Pole			1 pt.	1 qt.
	Large Lima	1 pt.	1 pt.	1 qt.	1 qt.
Beet ·····	Extra Early Turnip	1 oz. 1 oz.	2 oz. 2 oz.	4 oz. 8 oz.	8 oz.
	Long Blood	102.	2 02.	4 oz.	8 oz
Brocoli	Purple Cape			1/2 OZ.	12 OZ.
Brussels	Sprouts		7 4,4 4	1/2 OZ.	1/2 OZ.
Cabbage	Early Ox-Heart	½ oz.	16.00	16.00	2 OZ.
	Early Winningstadt	1/4 OZ.	4 0Z.	½ 0Z. ½ 0Z.	2 OZ.
	Large Flat Dutch	1/2 OZ.	1/2 OZ.	1 oz.	1 02
	Red Dutch for Pickling		1/8 OZ.	1/2 OZ.	1/2 OZ.
Carrot	Early Horn	1/2 OZ.	1/2 OZ.	1 oz	1 oz.
0	Long Orange	1 oz.	2 oz.	4 oz	4 oz.
Cauliflower	Nonpareil	% OZ.	4 OZ.	1/4 OZ 1/2 OZ.	2 OZ.
Celery	Early Dwart Sugar	1 pt.	1 qt.	2 qt.	% oz. 2 qt.
00/11	Eight Rowed Sugar		1 pt.	1 qt.	2 qt.
	Evergreen	½ pt.	1 pt.	1 qt.	1 qt.
Corn Salad.		1 00	1/2 OZ.	1 oz.	2 oz.
Cress, OR	Peppergrass	1 oz.	2 oz. 1 oz.	1 oz.	4 oz.
Cucumber	Long Green	/2 02.	102.	1/2 OZ.	1 oz. 1 oz.
Egg Plant	Improved New-York Purple, .	1/8 OZ.	. 1/4 OZ.	1/2 OZ.	1 oz.
	Scarlet Chinese				8 OZ.
Endive	Green Carled	1/4 OZ.	½ 0Z.	1 oz.	2 oz.
Kale Kohlrabi	Green Curled Scotch	1/4 OZ.	½ 0Z. ½ 0Z.	1 oz.	1 oz.
Leek	Best Flag	14 OZ.	1/4 OZ.	1 oz.	1 oz. 1 oz.
Lettuce	Early Curled Simpson's	1/4 OZ.	14 OZ.	½ OZ.	1 oz.
	Butter		1/4 OZ.	1/2 OZ.	1/2 OZ.
********	Ice Drumhead	1/4 OZ.	1/2 OZ.	1/2 OZ.	1/2 OZ.
B#I o loss	White Cabbage	14 OZ. 14 OZ.	½ 0Z. ½ 0Z.	½ oz. 1 oz.	1/2 OZ.
Melon	Nutmeg	74 02.	1/4 OZ.	1/2 OZ.	1 oz.
********	Ice Cream Water	½ 0Z.	1 oz.	2 oz.	2 oz.
Okra	Long Green	1 oz.	2 oz.	4 oz.	8 oz.
Onion	Yellow Danvers	1 oz.	1 oz.	2 oz.	4 oz.
******	White Portugal	½ 0Z.	1 oz. 1 oz.	2 oz. 2 oz.	4 oz.
Parsley	Extra Curled	13 021	1/2 OZ.	1 oz.	4 oz 1 oz.
Parsnip	Student	1 oz.	1 oz.	4 oz.	8 oz.
Peas	Carter's First Crop	1 pt.	1 qt.	2 qt.	2 qt.
*******	McLean's Advancer	½ pt.	1 pt.	1 qt.	2 qt.
	Harrison's Perfection Champion of England	i pt.	i qt.	1 qt. 2 qt.	2 qt.
**********	Eugenie	1 pt.	i qt.	1 qt.	4 qt. 2 qt.
*******	Hair's Mammoth			1 qt.	1 qt
Pepper	Sweet Mountain		1/4 OZ.	1/2 OZ.	1/2 OZ.
Padich	Large Squash	1 oz.	2 oz.	1/4 OZ.	1/4 OZ.
Radish	Scarlet Turnip Yellow Turnip	1 02,	2 0Z.	4 oz. 1 oz.	8 oz.
********	Long Scarlet	1 oz.	1 oz.	4 oz.	4 oz. 4 oz.
	Chinese Winter		1/4 OZ.	1/2 02.	1 oz.
Salsify Spinach Squash	Flandra	½ 0Z.	2 oz.	4 oz.	8 oz.
Squash	Flandre Early Bush Crook-Neck	1/4 lb. 1/2 oz.	1/4 lts.	1/2 lts.	1 lb.
oquasii	Hubbard	1/2 OZ.	1/2 OZ.	1 oz. 1 oz.	2 oz. 1 oz.
	Yokahama	1/4 OZ.	1/4 OZ.	½ oz.	1 oz.
Tomato	Early Red Smooth	1/4 OZ.	1/4 OZ.	1/2 OZ	1 0Z.
*****	Large Yellow		1/4 OZ.	1/2 OZ.	1/2 OZ.
Turnin	Filden	1/4 OZ.	1/4 OZ.	1/2 OZ.	1 oz.
Turnip	White French	1/2 OZ.	½ oz. 1 oz.	½ OZ. 2 OZ.	1 oz.
	Yellow Stone.	1 oz.	1 oz.	2 oz.	4 oz. 4 oz.
Herbs	Yellow Stone	****	1 pa ea.	1 pa. ea.	1 pa. ea.
				-	Lui sale

September.	SEPTEMBER.	Cabbage. Cabbage. Could Frames. Corn Salad. Corn Salad. Corn Salad. Corn Salad. Cress. Chres. Mistard. Winter Radish, Spinach, Turnip.
TABLE calling to mind the various Vegetable Seeds that may be sown in each month from February to September.	AUGUST.	sow in open ground.  Articholke, Asparagus, Brands, Grands Brands, Brands Brands, Grands Brands, Grands, G
1 each month fr	JULY.	sow in open groups. Beans, Beet, Sweet Corn, Sweet Corn, Corn Salad, Cress, Gucumber, Endive, Gherkin, Kale, Kollrabi, Lettuce, Nasturtium, Okra, Early Peas, Pumpkin, Early Peas, Pumpkin, Spinach, Spin
may be sown in	JUNE.	Bush Beans, Liuna do. Bush Beans, Liuna do. Bearle, Bearle, Beret, Beret, Beret, Beret, Beret, Brocoli, Brocoli
able Seeds that	MAY.	sow in open ground of a strict
e various Veget	APRIL.	sow is horner. Sweet Corn, Gucumber, Gucumber, Pepper, Tomato.  sow in ores geous English Beans, Bredgish Beans, Bredgish Beans, Bredgish Bears, Carliflower, Calery, Carliflower, Carliflo
ling to mind th	MARCH.	sow in hot-bed. Early Beaus, Early Beach, Brussels Sprouts, Brusels Sprouts, Cauliflower, Persiew, Persiew, Persiew, Persiew, Persiew, Persiew, Cauliflower, Cauliflower, Carrot, Car
TABLE ca	FEBRUARY.	Sow IN HOT NED.  Early Beans, Brotoli, Early Cabbage, Foreing Carrot, Early Celery, Cacumber, Early Lettuce, Early Lettuce, Paper, Paper, Radish, Tomato.